

Rituximab and hyaluronidase human (Rituxan Hycela)

Provider Order Form



Newport Superior Infusion

1501 Superior Ave, Suite 202

Newport Beach, CA 92663

Phone: 949-601-6001

Fax: 866-542-8631

PATIENT INFORMATION

Date:	Patient Name:	DOB:
ICD-10 code (required):		ICD-10 description:
☐ NKDA Allergies:	Weight lbs / kg:	Height in / cm:
Patient Status: ☐ New to Therapy ☐ Continuing Therapy Next Due Date (if applicable):		

PROVIDER INFORMATION

Ordering Provider:	Provider NPI:		
Referring Practice Name:	Phone:	Fax:	
Practice Address:	City:	State:	Zip Code:

NURSING

- ☒ Provide nursing care per Newport Infusion Standard Procedures
- ☐ Hepatitis B status and date (list results and attach clinicals):

RECOMMENDED PRE-MEDICATION ORDERS

- ☐ acetaminophen (Tylenol) ☐ 500mg / ☐ 650mg / ☐ 1000mg PO
- ☐ diphenhydramine (Benadryl) ☐ 25mg / ☐ 50mg ☐ PO / ☐ IV
- ☐ methylprednisolone (Solu-Medrol) 125mg IV

PRE-MEDICATION ORDERS

- ☐ cetirizine (Zyrtec) 10mg PO
- ☐ loratadine (Claritin) 10mg PO
- ☐ famotidine (Pepcid) 20mg PO
- ☐ Other: _____
Dose: _____ Route: _____
Frequency: _____

LABORATORY ORDERS

- ☐ CBC ☐ at each dose ☐ every _____
- ☐ CMP ☐ at each dose ☐ every _____
- ☐ CRP ☐ at each dose ☐ every _____
- ☐ Other: _____

SPECIAL INSTRUCTIONS

THERAPY ADMINISTRATION

- ☒ **Rituximab and hyaluronidase human (Rituxan Hycela)** subcutaneous injection
 - Dose
 - ☐ 1400mg / 23,400 units per 11.7mL
 - ☐ 1600mg / 26,800 units per 13.4mL
 - ☐ Other: _____
 - Frequency
 - ☐ Administer once weekly for 3 weeks following full dose of rituximab product by intravenous infusion on week 1 (4 weeks total)
 - ☐ Administer once weekly for 7 weeks following full dose of rituximab product by intravenous infusion on week 1 (8 weeks total)
 - ☐ Administer on day 1 of Cycles 2-8 (every 21 days) for up to 7 cycles following a full dose of a rituximab product
 - ☐ Following completion of 6-8 cycles of CVP chemotherapy and a full dose of rituximab product intravenous infusion at week 1, administer once weekly for 3 weeks at 6 month intervals to a maximum of 16 doses
 - ☐ Administer on day 1 of Cycles 2-6 for a total of 5 cycles following intravenous dose at day 1
 - ☐ Other: _____
 - Rate
 - ☐ Administer first infusion over 90 minutes
 - ☐ Administer subsequent infusions over 30 minutes if previous
- Refills: ☐ Zero / ☐ for 12
☐ _____ (if not indicated order will expire one year from date signed)

Provider Name (Print)

Date

Provider Signature